



## Maisflocken

### Enkelvoudig voeder

### Enkelvoudig voeder

De voordelen zijn onder meer het hogere energiegehalte in vergelijking met haver met een kleinere hoeveelheid eiwit. Een onbetwist nadeel van maïs is echter de harde structuur, wat betekent dat er slecht op wordt gekauwd en dat de waardevolle ingrediënten soms onvoldoende worden verteerd. Als maïs echter hydrothermisch met vochtige hitte wordt ontsloten, ontwikkelen de resulterende maïsvlokken hun voordelen bij het voederen van paarden: De vlokvorming verhoogt de verteerbaarheid en tolerantie van maïs aanzienlijk en maakt het veel smakelijker. Gebruik maïsvlokken om het gebruikelijke kribvoederrantsoen op te waarderen of als substantiële basis voor een individueel samengesteld kribvoeder. Trouwens: 0,8 kg maïs vervangt ongeveer 1 kg haver in energiewaarde.

### Aanbevolen voeder:

### Feeding recommendation:

for horses:

- Due to the high energy density in connection with the content of valuable fatty acids and carotenoids, corn flakes are often fed to sport and breeding horses to upgrade their feed diet in addition to sufficient roughage.
- As an alternative to oats, 0.8kg of corn flakes replace 1kg of oats.
- In case of liver insufficiency and kidney diseases with high urea values in the blood plasma, Corn Flakes are recommended as an energy supply in protein-reduced diets.

for chicken and other poultry species:

- For laying hens and breeders, Corn Flakes are working as an energetic upgrade in the daily feed mixture and complete the diet with high content of essential fatty acids and carotenoids.
- This supports a well pigmentation of the egg yolk.
- In pullet feed, we recommend limiting the mixing rate of Corn Flakes to a maximum of 30% in order to prevent unwanted fat formation and increased fat deposits in the ovaries
- We also recommend a maximum of 30% Corn Flakes in the feed rations for waterfowl in order to avoid disadvantageous fat formation.

for small animals:

- Corn flakes should only be fed to rabbits and rodents in moderation or in phases with increased energy requirements, such as to feed weakened animals.
- Because the digestive tract of our little darlings is not designed for a high





starch and energy content in the feed, which is why excessive grain feeding can quickly lead to obesity and metabolic problems.

- To stimulate the activity and the search for food, we recommend mixing corn flakes with 1 teaspoon once or twice a week into the feed ration or “hiding” them in the hay/fresh bedding.

For the diet construction, don't hesitate to contact us for individual advices which we recommend to closely coordinate with the treating vet.

verteerbaar Eiwit (vRp): 59,9 g/kg

prececal digestible protein (pcvRp): 54,7 g/kg

verteerbare Energie (MJ DE): 13,6 MJ DE/kg

Metabolizable energy (MJ ME): 12,8 MJ ME/kg

**Analytische componenten en gehaltes:** 8,50 % Ruw eiwit, 3,00 % Ruw vet, 1,80 % Ruwe vezels, 1,00 % Ruwe as, 0,03 % Calcium, 0,30 % Fosfor, 61,60 % Zetmeel, 1,80 % Suiker

